

Studying the stellar populations in the fuzzy cores of young massive clusters

Zeinab Khorrami

P. Clark (Cardiff University), F. Vakili (Observatoire de la Cote d'Azur),
T. Lanz (Observatoire de la Cote d'Azur)

29 June 2017

Cardiff University



Young Massive clusters

Dense aggregate of newly formed stars with lots of difficulties!


- Immersed in gas/dust: High (inhomogeneous) extinction
Observations in longer wavelength
- Most of them are not close enough: their individual members probably are not resolved
High angular resolution observations
- Massive hot stars mask the faint low-mass stars.
High contrast imaging

DSS2 Blue




FoV: 23.03'

DSS2 Red




FoV: 23.03'

2MASS J,H,K



FoV: 23.03'

2MASS J,H,



Next step ...

1. Proper evolutionary/atmosphere model
2. Estimating stellar masses, ages
3. Investigate physics of the newly formed cluster:
 - MF, density profile, Virial status, mass-segregation
 - Feeding cluster-formation simulations, gravitational-dynamical evolution
 - Filling the gap between cluster formation and evolution

Final step ...

To compare simulations with observations we should use
synthetic observations

With **MYSO** feel free to

Make Your Synthetic Observations!

Two similar clusters (age and stellar population)
in different distances


NGC 3603




LMC


SMC

NGC 3603



NGC 3603





$M_{total} \sim 10^4 M_{\odot}$

$Age \sim 1 - 2 Myr$

$Dis \sim 6 - 7 Kpc$


Constellation : Carina

NGC 3603

| MF slope | condition | Reference |
|------------------|-------------------------|-------------------------|
| -0.73 | $(1 - 30) M_{\odot}$ | [Eisenhauer et al.1998] |
| -0.9 | $(2.5 - 100) M_{\odot}$ | [Sung & Bessell2004] |
| -0.5 ± 0.1 | $r < 6''$ | [Sung & Bessell2004] |
| -0.8 ± 0.2 | $6'' - 12''$ | [Sung & Bessell2004] |
| -1.2 ± 0.2 | $r > 12''$ | [Sung & Bessell2004] |
| -0.91 ± 0.15 | $(0.4 - 20) M_{\odot}$ | [Stolte et al.2006] |
| -0.31 | $0 - 5''$ | [Harayama et al.2008] |
| -0.55 | $5'' - 10''$ | [Harayama et al.2008] |
| -0.72 | $10'' - 13''$ | [Harayama et al.2008] |
| -0.75 | $13'' - 30''$ | [Harayama et al.2008] |
| -0.26 | $0 - 5''$ | [Pang et al.2013] |
| -0.55 | $5'' - 10''$ | [Pang et al.2013] |
| -0.76 | $10'' - 15''$ | [Pang et al.2013] |


Different observations


Different MF slopes




WFPC2
Filters: V,R,I
50mas/pix

ACS/HRC
Filters: Ux,U,B,V,R,I
25mas/pix







WFPC2 shows the decreasing trend in MF slope... signature of mass-segregation BUT this is not the case for HRC data



High angular and contrast images from VLT/SPHERE

MF example: NGC3603





No signature of mass segregation in the core of NGC 3603:


- 1) The MF slope in its very core is not flatter than the next radial bin
- 2) Both slopes are similar to the MF values found in previous works for the outer regions


[Khorrami et al., 2017, A&A, 588, id.L7]

R136









30 Doradus Nebula and Star Cluster

Hubble Space Telescope • WFC3

Visible WFC3/UVIS

F336W *U* F438W *B* F555W *V* F814W *I* F656N H α

50 light-years
15.3 parsecs 61''


F110W *J* F160W *H*





Dis \sim 50 Kpc
Constellation : 30 Doradus

$M_{total} \sim 10^5 M_\odot$
Age $\sim 2 - 3 Myr$


Mass function slopes for R 136 from previous analyses.

| MF slope | Condition | Reference |
|------------------|---|------------------------|
| -0.90 | (20–70) M_{\odot} $r < 3'.3$ | Malumuth & Heap (1994) |
| -1.89 | (20–70) M_{\odot} $3'.3 < r < 17'.5$ | Malumuth & Heap (1994) |
| -1.0 ± 0.1 | (2.8–15) M_{\odot} $2'.0 < r < 18'.8$ | Hunter et al. (1996) |
| (-1.3)–(-1.4) | (15–120) M_{\odot} | Massey & Hunter (1998) |
| -1.59 | $r < 1'.6$ | Brandl et al. (1996) |
| -1.33 | $1'.6 < r < 3'.2$ | Brandl et al. (1996) |
| -1.63 | $3'.2 < r$ | Brandl et al. (1996) |
| -1.17 ± 0.05 | $4''.6 < r < 19''.2$ | Selman et al. (1999) |
| -1.37 ± 0.08 | $15'' < r < 75''$ | Selman et al. (1999) |
| -1.28 ± 0.05 | (2–6.5) M_{\odot} $4'' \lesssim r \lesssim 20''$ | Sirianni et al. (2000) |
| -1.2 ± 0.2 | (1.1–20) M_{\odot} $20'' < r < 28''$ | Andersen et al. (2009) |





MF example: R136



[Khorrami et al., 2017, A&A, 602, A56]

MF is sensitive to the observations
and models

How can we simulate observations?

Make Your Synthetic Observations

Inputs:

Stars information (from N-body): 3D position and Velocity, Mass, age, metallicity

Cloud information (from SPH): 3D position, particle's mass, smoothing lengths

Observational Filter (from the list)

Imaging angular resolution AND Spectroscopic resolution

Distance of the centre of mass

FoV

R_v for extinction


Adaptive optics: seeing and SR

OB-treatment: TLUSTY model atmosphere

Velocity dispersion


Euler angles for line-of-sight

Signal/Noise ratio for the faintest star





***_cube_spectra.fits** : 3D cube,
X-Y is the position of stellar
sources, z is flux in different
wavelengths

Gaia-G filter
R=700





*_cube_spectra.fits : 3D cube,
X-Y is the position of stellar
sources, z is flux in different
wavelengths

Gaia-G filter
R=700




*_cube_spectra.fits : 3D cube,
X-Y is the position of stellar
sources, z is flux in different
wavelengths


Gaia-G filter
R=700



$M_{\text{tot}} = 10^5 M_{\odot}$, BF=0% , $R_h = 0.8 \text{ pc}$, Age=2 Myr




WFPC2/F555W



IRDIS/K
SR: 0.7

Synthetic Observations with MYSO

 $M_{\text{tot}} = 10^5 M_{\odot}$, BF=0%, $R_h = 0.8 \text{ pc}$, Age=2 Myr


IRDIS/Ks: SR 0.7

SR 0.8




SR 0.9


HST/WFC3/F555W

Non-segregated






Segregated







SR 0.9


HST/WFC3/F555W



MF slope
 $K \sim -1.38$
 $V \sim -0.93$



Minimum Spanning Tree




$$\Lambda_{\text{MSR}} = \frac{\langle l_{\text{norm}} \rangle}{l_{\text{massive}}} \pm \frac{\sigma_{\text{norm}}}{l_{\text{massive}}}$$



$\Lambda_{\text{MSR}} = 1.0$: No-segregation

$\Lambda_{\text{MSR}} > 1.0$: Segregated


[Allison et al. 2009]



FoV: 16"x16"
VLT/SPHERE/K
SR=0.75
Seeing=0.8"



FoV: 16"x16"
HST/WFPC2/F555W




MST method for mass-segregation


FoV: 16" x 16"
VLT/SPHERE/K
SR=0.75
Seeing=0.8"

$M_{\text{tot}} = 10^4 M_{\odot}$, BF=50%, $R_h = 0.5 \text{ pc}$, D=3.0



Segregated, SPHERE/IRDIS/K
Segregated, HST/WFPC2/F555W
Not-segregated, SPHERE/IRDIS/K
Not-segregated, HST/WFPC2/F555W




FoV: 16" x 16"
HST/WFPC2/F555W




FoV: 16"x16"
VLT/SPHERE/K
SR=0.75
Seeing=0.8"



FoV: 16"x16"
HST/WFPC2/F555W



MST method for mass-segregation




Summary and conclusion

- MF is sensitive to the resolution of the observational instrument
 - Observers need to compare the data with different resolution
 - We always need higher angular resolution data with better contrast
- Synthetic observations are needed to compare simulations with

Observations

- MST method can detect mass-segregation in the simulated data



A nighttime photograph of the Prague skyline. In the background, the Gothic St. Vitus Cathedral with its multiple spires and intricate stonework is prominent. In the foreground, the Charles Bridge (Karluv Most) spans the Vltava River, its arches reflected in the dark water. The city buildings are illuminated by streetlights and building lights, creating a warm glow against the night sky.

Thanks for your attention!

Děkujeme za pozornost!